

BAB V **PENUTUP**

5.1 Kesimpulan

Penilaian keanekaragaman dan status regenerasi dari spesies pohon adalah penting untuk pemanfaatan, pengelolaan, dan pelestarian berkelanjutan. Dalam penelitian ini, struktur populasi dari seluruh spesies pohon mengungkapkan bahwa kontribusi anakan terhadap total populasi adalah yang tertinggi diikuti oleh dewasa dan semai. Fakta ini menunjukkan bahwa status regenerasi secara keseluruhan dari spesies pohon di TAHURA Buraen adalah "adil" dan "baru" komunitas di masa depan dapat dipertahankan kecuali ada tekanan lingkungan utama atau gangguan yang ditimbulkan oleh manusia. Akan tetapi, pertumbuhan, kelangsungan hidup, dan potensi reproduksi spesies pohon yang menunjukkan regenerasi "buruk" atau "tidak ada" regenerasi, mungkin akan berisiko di masa depan. Oleh karena itu, perencanaan pengelolaan (manajemen) yang sistematis diperlukan untuk konservasi dan pemanfaatan sumber daya tumbuhan secara berkelanjutan.

5.2 Saran

Saran yang dapat diberikan dalam penelitian ini adalah:

- 1) Dibutuhkan perhitungan dan analisis tentang disparitas antara habitat pinggir hutan dan habitat interior hutan
- 2) Ruang sempling dapat diperpanjang hingga 1 km ke dalam hutan untuk melihat perbandingan yang lebih luas dan komprehensif.

DAFTAR PUSTAKA

Sumber acuan dari Buku dan Journal:

- Alam Nasional Indonesia.2008.Bakossutraman(Badan Koordinasi Survey dan Pemetaan Nasional).Cibinong,Bogor.
- Alfian, M. 2014. Keanekaragaman Hayati. Jakarta: Universitas Islam Negeri Syarif Hidayatullah
- Backer, C.A. and van den Brink, R.C.B. 1963. *Flora of Java*. N.V.P. Nordhoff, Groningen, The Netherlands.
- Baduni NP, Sharma CM. 1999. Community structure and growing stock variation in *Quercus floribunda* forest on different aspects of Garhwal Himalaya. Bangladesh Journal of Forest Science 28:82-93.
- Banilodu L. and Saka N.T. 1993. Descriptive Analysis of Sumba Forests. Widya Mandira Catholic University. Kupang.
- Ballabha R, Tiwari JK, Tiwari P. 2013. Regeneration of tree species in the sub-tropical forest of Alaknanda Valley, Garhwal Himalaya, India. Forest Science and Practice 15:89-97.
- Clark JS, Beckage B, Camill P, et al. 1999. Interpreting recruitment limitation in forests. American Journal of Botany 86:1-16.
- Dalling JW, Hubbel SP, Silver K. 1998. Seed dispersal, seedling establishment and gap partitioning among tropical pioneer trees. Journal of Ecology 86:674-689.
- Drees, E.M. 1950. *Daftar Nama-nama Pohon-pohon Pulau Timor*. Laporan Balai Penyelidikan Kehutanan. Nomor 33. Balai Penyelidikan Kehutanan. Bogor, Indonesia.
- Eilu G, Obua J. 2005. Tree condition and natural regeneration in disturbed sites of Bwindi impenetrable forest National Park, Southwestern Uganda. *Tropical Ecology* 46:99-111.
- Elouard C, Pascal JP, Pelissier R, et al. 1997. Monitoring the structure and dynamics of a dense moist evergreen forest in the Western Ghats (Kodagu District, Karnataka, India). *Tropical Ecology* 38:193-214.

Flora of Indonesia. https://en.wikipedia.org/wiki/Flora_of_Indonesia [Last accessed 15 May 2018].

Grubb PJ. 1977. The maintenance of species richness in plant communities. The importance of the regeneration niche. *Biological Reviews* 52:107-145.

Gairola S. 2010. Phytodiversity (angiosperms and gymnosperms) in Mandal-Chopta forest of Garhwal Himalaya, Uttarakhand, India. PhD thesis. Garhwal, Uttarakhand: HNB Garhwal University Srinagar.

Ghildiyal S, Baduni NP, Khanduri VP, et al. 1998. Community structure and composition of oak forests along altitudinal gradient in Garhwal Himalaya. *Indian Journal of Forestry* 21:242-247.

Jones RH, Sharitz RR, Dixon PM, et al. 1994. Woody plant regeneration in four flood plants forests. *Ecological Monographs* 64:345-367.

Khumbongmayum AD, Khan ML, Tripathi RS. 2005. Survival and growth of seedlings of a few tree species in the four sacred groves of Manipur, Northeast India. *Current Science* 88:1781-1788.

Krauchi N, Brang P, Schonenberger W. 2000. Forests of mountainous regions: gaps in knowledge and research needs. *Forest Ecology and Management* 132: 73-82.

Millennium Ecosystem Assessment (MA). 2005. *Ecosystems and Human Well-Being: Synthesis*. Island Press, Washington, DC.

Malik ZA. 2014. *Phytosociological behaviour, anthropogenic disturbances and regeneration status along an altitudinal gradient in Kedarnath Wildlife Sanctuary (KWLS) and its adjoining areas*. PhD thesis. Uttarakhand: HNB Garhwal University Srinagar Garhwal.

Malik ZA, Pandey R, Bhatt AB. 2016. Anthropogenic disturbances and their impact on vegetation in Western Himalaya, India. *Journal of Mountain Science* 13:69-82.

Malik ZA, Bhatt AB. 2015. Phytosociological analysis of woody species in Kedarnath Wildlife Sanctuary and its adjoining areas in Western Himalaya, India. *Journal of Forest and Environmental Science* 31:149-163.

Malik ZA, Bhatt AB. 2016. Regeneration status of tree species and survival of their seedlings in Kedarnath Wildlife Sanctuary and its adjoining areas in Western Himalaya, India. *Tropical Ecology*, in press.

- Mishra AK, Behera SK, Singh K, et al. 2013. Influence of abiotic factors on community structure of understory vegetation in moist deciduous forests of north India. *Forest Science and Practice* 15:261-273.
- Odum EP. 1971. Fundamentals of ecology. Philadelphia (PA): Saunders.
- Peet RK. 1974. The measurement of species diversity. *Annual Review of Ecology, Evolution, and Systematics* 5:285-307.
- Raturi GP. 2012. Forest community structure along an altitudinal gradient of district Rudraprayag of Garhwal Himalaya, India. *Ecologia* 3:76-84.
- Rushayati, S.B. 2007. Mengenal Keanekaragaman Hayati. Yayasan Kehati: Grasindo
- Saberwal VK. 1995. Pastoral politics: Gaddi grazing, degradation and biodiversity conservation in Himachal Pradesh, India. *Conservation Biology* 10
- Sarkar M, Devi A. 2014. Assessment of diversity, population structure and regeneration status of tree species in Hollongapar Gibbon Wildlife Sanctuary, Assam, Northeast India. *Tropical Plant Research* 1:26-36.
- Saikia P, Khan ML. 2013. Population structure and regeneration status of Aquilaria malaccensis Lam. in home gardens of Upper Assam, northeast India. *Tropical Ecology* 54:1-13.
- Simpson EH. 1949. *Measurement of diversity*. Nature 163:688.
- Sharma CM, Mishra AK, Prakash O, et al. 2014. Assessment of forest structure and woody plant regeneration on ridge tops at upper Bhagirathi Basin in Garhwal Himalaya. *Tropical Plant Research* 1:62-71.
- Shaheen H, Ullah Z, Khan SM, et al. 2012. Species composition and community structure of western Himalayan moist temperate forests in Kashmir. *Forest Ecology and Management* 278:138-145.
- Saxena AK, Singh JS. 1984. Tree population structure of certain Himalayan forest associations and implications concerning the future composition. *Vegetatio* 58: 61-69.
- Tewari JC, Singh SP. 1985. Analysis of woody vegetation in a mixed oak forest of Kumaon Himalaya. *Proceedings of Indian National Science Academy* 51:332-347.
- Verma RK, Kapoor KS, Subramani SP, et al. 2004. Evaluation of plant diversity and soil quality under plantation raised in surface mined areas. *Indian Journal of Forestry* 27:227-233.

Whittaker RH. 1965. Dominance and diversity in land plant communities. Science 147:250-260

Zobel DB, McKee A, Hawk GM, et al. 1976. Relationships of environment to the composition, structure and diversity of forest communities of the central western cascades of Oregon. Ecological Monographs 46:135-156.

Sumber Acuan dari Website Link:

<http://bioimages.vanderbilt.edu/tree-key/index.htm> [last accessed August 16, 2018]